

Sampling rare events: statistics of local sequence alignments

Alexander K. Hartmann

Department of Physics, University of California, Santa Cruz CA 95064, USA

E-mail: hartmann@bach.ucsc.edu

(February 1, 2008)

Summary A new method to simulate probability distributions in regions where the events are *very unlikely* (e.g. $p \sim 10^{-40}$) is presented. The basic idea is to represent the underlying probability space by the phase space of a physical system. The system is held at a temperature T , which is chosen such that the system preferably generates configurations which originally have low probabilities. Since the distribution of such a physical system is known from statistical physics, the original unbiased distribution can be obtained.

As an application, local alignment of protein sequences based on BLOSUM62 substitution scores with (12,1) affine gap costs are considered. The distribution of optimum sequence-alignment scores S is studied numerically over a large range of scores. The deviation of $p(S)$ from the extreme-value (or Gumbel) distribution is quantified. This deviation decreases with growing sequence length.

Introduction Modern molecular biology, e.g. the *human genome project* [1], relies heavily on the use of large databases [2,3], where DNA or protein sequences are stored. The basic tool for accessing these databases and comparing different sequences is *sequence alignment*. The result of each comparison is an maximum alignment *score* S . One is interested either in *global* or *local* optimum alignments. For the first case, the score is maximized over all alignments which include both sequences fully, while for the second case, the optimum is calculated over all possible subsequences. To estimate the significance of the result of a comparison, one has to know, based on a random model, the statistical distribution $p(S)$ of scores. For biologically relevant models, e.g. for protein sequences with BLOSUM62 substitution scores [4] and affine gap costs [5], $p(S)$ is not known in the interesting region, where $p(S)$ is small. A number of empirical studies [6–10] for local alignment, in the region where $p(S)$ is large, suggest that $p(S)$ is an *extreme value* (or Gumbel) distribution [11]

$$p_G(S) = \lambda e^{-\lambda(S-u)} \exp(-e^{-\lambda(S-u)}), \quad (1)$$

where u denotes the maximum of the distribution and λ characterizes the behavior for large values of S , i.e. the tail of the distribution.

In this work, to determine the tail of $p(S)$, a *rare event simulation* is applied. For dynamical problems, like investigating queuing systems or studying the reliability of technical components, several techniques [12], mostly based on *importance sampling* have been developed. Related methods have been introduced in physics, like *multi-canonical sampling* [13] or the *pruned-enriched Rosenbluth method* [14].

By simply changing perspective, one can apply these standard techniques to many other problems. Here, the method it is applied to the sequence alignment problem. Instead of directly drawing the random sequences, the basic idea is to use a physical system, which has a state given by a pair of sequences and is held at temperature T . This is similar to the simulated annealing approach [15], used to solve hard optimizations problems. The state of the system changes in time, governed by the rules of statistical mechanics. The energy E of the system is defined as $E = -S$. Therefore, at low temperatures the system prefers a state with high score value S . Since the thermodynamic properties of the system are known, it is possible to extract from the measured distribution $p^*(S)$ of scores the target distribution $p(S)$.

For the alignment problem two sequences $\mathbf{x} = x_1 x_2 \dots x_n$ and $\mathbf{y} = y_1 y_2 \dots y_m$ over a finite alphabet with r letters are given. For DNA the alphabet has 4 letters, representing the bases, for protein sequences it has 20 letters, representing the amino acids. Let f_i be the probability for the occurrence of letter i , assuming here that all letters of a sequence are independent. An alignment is a pairing $\{(x_{i_k}, y_{j_k})\}$ ($k = 1, 2, \dots, K$, $1 \leq i_k < i_{k+1} \leq n$ and $1 \leq j_k < j_{k+1} \leq m$) of letters from the two sequences. Please note that some letters may not be aligned, i.e. *gaps* occur. To each alignment a score is assigned, via a scoring function $s(x, y)$. The total score is the sum of scores of all aligned letters $\sum_k s(x_{i_k}, y_{j_k})$ plus the costs of all gaps. Here, so called *affine* gap costs (α, β) are considered, i.e. a gap of length l has costs $g(l) = -\alpha - \beta(l - 1)$. For both global and local alignment efficient algorithms [16,17,5] exist, which calculate an optimum alignment in time $O(nm)$. Hence, one can easily generate e.g. $N \approx 10^5$ samples of pair of sequences according the frequencies f_i , obtain each time the optimum alignment, and calculate a histogram of the scores S . This *simple sampling* allows one to calculate $p(S)$ in the region where the probabilities are large (e.g. $p(S) \approx 10^{-4}$). Recently, the *island method* [18] was introduced, which allows a speed up of several orders of magnitudes, but still the far end of the distribution is out of reach.

Algorithm The basic idea to determine the behavior of $p(S)$ at the rare event tail (e.g. $p(S) \approx 10^{-40}$) is to view a pair of sequences as a physical system, which behaves according the rules of statistical mechanics. This means that instead of considering many independent pairs of fixed sequences, one pair of sequences $c(t) = (\mathbf{x}(t), \mathbf{y}(t))$ is used, which changes at discrete

times t . Mathematically speaking, a Markov chain [19] $c(0) \rightarrow c(1) \rightarrow c(2) \rightarrow \dots$ is used to generate the instances. The behavior of a Markov chain is not deterministic, but governed by probabilities $p(c \rightarrow c')$ that state c' follows immediately after state c .

The simplest rule for the transition is, to choose randomly a position in one of the sequences and choose randomly a new letter from the alphabet, with all positions being equiprobable and the letters having probabilities f_i , i.e. $p(c \rightarrow c') = f_i/(n+m)$ if c, c' differ by at most one letter, and $p(c \rightarrow c') = 0$ otherwise. With this choice of the transition probabilities, for $t \rightarrow \infty$ all possible pairs of sequence have the probability $P(c) = \prod_i f_{x_i} \prod_j f_{y_j}$ of occurring. Hence, simple sampling is reproduced.

To increase the efficiency, one can apply *importance sampling* [12,20], a standard method for simulating rare events [21], which allows one to concentrate the sampling in small regions in configuration space. Suppose, we want to calculate the expectation value $\langle g \rangle_P$ of a function g on a (here discrete) configuration space $\{c\}$, with respect to the probabilities $P(c)$. Thus, we have $\langle g \rangle = \sum_c g(c)P(c)$. To estimate $\langle g \rangle_P$, we generate N samples $c^{(i)}$ which are distributed according to P , and calculate $1/N \sum_i g(c^{(i)})$. Let now $Q(c)$ be another probability distribution with $Q(c) \neq 0$ for all c . Then we can write $\langle g \rangle_P = \sum_c \frac{g(c)P(c)}{Q(c)}Q(c) = \langle gP/Q \rangle_Q$. This means, when we generate N configurations $c^{(i)}$ according to the probabilities $Q(c)$, then $1/N \sum_i g(c^{(i)})P(c^{(i)})/Q(c^{(i)})$ is also an estimator for $\langle g \rangle_P$. By choosing Q such that the sampling occurs in the region of the phase space we are interested in, we can obtain a much higher accuracy there.

A good choice for the transition rule of the sequence-alignment problem is to select one position in the pair c of sequences randomly, change the letter randomly according the frequencies f_i , recalculate the optimum alignment $S(c')$ with a standard algorithm and accept this move $c \rightarrow c'$ with probability $\max(1, \exp(\Delta S/T))$, where $\Delta S = S(c') - S(c)$. This transition probabilities describe a physical system at temperature T with energy $E = -S$. The advantage of this approach is that the equilibrium distribution $Q(c)$ is known from statistical physics [22]: $Q(c) = P(c) \exp(S(c)/T)/Z$ with $Z(T) = \sum_c P(c) \exp(S(c)/T)$ being a normalization constant, called the partition function. Thus, the probability to have score S is

$$p^*(S) = \frac{\exp(S/T)}{Z(T)} \sum_c' P(c), \quad (2)$$

where the sum $\sum_c'$ runs over all sequences with score S . Thus, to obtain $p(S)$, one simulates the system at temperature T and records a histogram of scores $p^*(S)$. Then, the unbiased estimator for the distribution is $p(S) = \sum_c' P(c) = p^*(S)Z(T) \exp(-S/T)$. Note that $Z(T)$ is unknown a priori, but can be determined very easily, as shown below.

To describe the behavior of $p(S)$ over a wide range, the model must be simulated at several temperatures. For this reason, and because at low temperatures the dynamics are slow in general so we want to increase the efficiency, the model is simulated via the *parallel tempering* method [23,24]. Using this technique, the system is simulated at N_T different temperatures $T_0 < T_1 < \dots < T_{N_T}$ in parallel, i.e. with N_T independent pairs $c(T_i)$ of sequences. The main idea of parallel tempering is, that from time to time the configurations between neighboring temperatures T_i, T_{i+1} are exchanged according a probabilistic rule [23,24]. Here, each simulation step consists of one Markov step for each configuration c and one exchange step between one neighboring pair $c(T_i), c(T_{i+1})$.

Example Next, a simple example is given, illustrating how the method works. Optimum local alignments without gaps for sequences of equal length $m = n = 20$ and $r = 4$ letters, all having the same probability $1/4$, are calculated. For the test the following score is applied: $s(x, y) = 1$ if $x = y$ and $s(x, y) = -3$ otherwise. Two types of runs are performed: (a) initially, all pairs of sequences are random, and (b) each pair consists of two equal sequences. Thus, for the first type, initially the score is low, while for the second type the score is initially maximal. This provides a criterion for equilibration: if the average score for both types of initial configuration agree within error bars (at time t_0), the simulation is long enough. In Fig. 1 the average optimum score S from 1000 independent runs and four different temperatures T is shown.




FIG. 1. Average alignment score S as a function of step t for $n, m = 20$, 4 letters, local alignment without gaps for different temperatures T . For each temperature, 1000 simulations were started with two random sequences (low scores) and 1000 simulations with two equal sequences (high scores).

To obtain uncorrelated samples, only samples at t_0 ,

$t_0 + \tau$, $t_0 + 2\tau$ etc are taken, where τ is the characteristic time in which the score-score correlation

$$c_S(t_0, t) = \frac{\langle S(t_0)S(t) \rangle - \langle S \rangle^2}{\langle S^2 \rangle - \langle S \rangle^2} \quad (3)$$

decreases to $1/e$.

In Fig. 2 the raw distributions of S for two temperatures is shown together with a distribution from a simple sampling of $N = 10^4$ realizations. Clearly, with the statistical mechanics approach, the region of high scores is sampled much more frequently.




FIG. 2. Raw distribution of alignment scores S for the direct simulation and for $T = 0.57$ and $T = 0.69$ with $n = m = 20$, 4 letters and local alignment without gaps.

For low scores, the final distributions obtained from the simple sampling and from the finite-temperature simulation must agree. This can be used to determine the constant $Z(T)$. It is chosen such that the difference in an interval $[S_1, S_2]$ between the simple sampling distribution and the rescaled distribution at T is minimal. In the same way the constants at lower temperatures can be obtained by matching to distributions obtained before at higher temperatures. The final distribution is shown in Fig. 3. For each datapoint, the distribution with the highest accuracy was taken. For comparison, a simple sampling distribution obtained using a huge number of samples ($N = 10^9$) is shown. Both results agree very well. Please note that the distribution from the finite- T approach spans almost the entire interval $[0, 20]$. In principal, the region for very small score S can be investigated also using the new method by simulating at *negative* temperature. How powerful the new method is can be seen by looking at the right border of the interval, where a value $p(20) = 9.13(20) \times 10^{-13}$ was obtained. This agrees within error bars with the exact result [25]

$0.25^{20} \approx 9.09 \times 10^{-13}$. This example illustrates, that the method presented here is indeed able to calculate accurately the distribution $p(S)$ of optimum alignment scores in regions where $p(S)$ is very small.




FIG. 3. Rescaled distribution $p(S)$ for the direct simulation and for $T = 0.57$, $T = 0.69$ for $n, m = 20$, 4 letters, local alignment without gaps. The solid line is the result of a large simple sampling simulation with $N = 10^9$ samples.

Results Next, the results for a biologically relevant case are presented. Sequences of amino acids distributed according the background frequencies by Robinson and Robinson [26] are used together with the BLOSUM62 scoring matrix [4] for $(12, 1)$ affine gap costs. This type of system has been studied in Ref. [10] in the region where $p(S)$ is large. Here, sequences of length $n = m$ in the range $[40, 400]$ were considered. The simulations were performed for $n_T = 7$ temperatures $T \in [2 \dots 10]$ ($[3.5 \dots 10]$ for $n, m = 400$), with up to 100 independent runs of length up to $t_{\max} = 4 \times 10^5$ steps. For the lowest temperatures it was not possible to equilibrate the system within the given time. The reason is that near $T \approx 1/\lambda$ the equilibration times seem to diverge. This indicates a phase transition in the physical system with (probably) a glassy phase at low temperatures. Hence, for the evaluation the only temperatures used, were those where equilibration was possible.

In Fig. 4 the distributions $p(S)$ of optimum alignment scores obtained in this way are shown. It was possible to determine the distribution in regions where $p(S)$ is as small as 10^{-40} . To obtain the same accuracy with a simple-sampling approach, given a computer which optimizes say 10^6 samples per second, a total simulation time of about 2.5×10^{17} times the age of the universe would be necessary. Also shown in Fig. 4 are fits of the low-score data to Gumbel distributions. The resulting parameters (λ, u) are comparable to the values found [10] before and depend slightly on the sequence length. For high scores,




FIG. 4. Distribution of alignment scores S for $L = 40, 100, 200, 400$, BLOSUM62 substitution matrix and affine (12,1) gap costs. The thin solid lines are fits to extreme-value distributions with parameters (λ, u) , yielding $(\lambda, u) = (0.355(5), 15.35(4))$ ($n, m = 40$), $(0.304(2), 21.67(4))$ ($n, m = 100$) and $(0.280(3), 32.01(3))$ for $n, m = 400$.

significant deviations are visible, in contrast to the earlier predictions. Since the deviations occur at high score values, they could not be detected before using conventional methods. The reason for the deviations is edge effects: very long alignments cannot start near the end of either of the sequences, so they become even more unlikely. The results found here can be fitted very well to *modified* Gumbel distributions of the form

$$\tilde{p}_G(S) = k\lambda e^{-\lambda(S-u)-\lambda_2(S-u)^2} \exp(-e^{-\lambda(S-u)}), \quad (4)$$

with $k \approx 1$, resulting in values for (λ, λ_2, u) of $(0.3277 \pm 0.0003, 8.56 \times 10^{-4} \pm 3 \times 10^{-6}, 15.35 \pm 0.04)$ for $n, m = 40$, $(0.2783 \pm 0.0003, 1.72 \times 10^{-4} \pm 1 \times 10^{-6}, 21.67 \pm 0.04)$ for $n, m = 100$, and $(0.2733 \pm 0.0004, 6.1 \times 10^{-5} \pm 2 \times 10^{-6}, 32.01 \pm 0.03)$ for $n, m = 400$. Anyway, with increasing lengths n, m , on a scale of scores proportional to $u \sim \log n$, $p(S)$ approaches the Gumbel distribution more and more, i.e. $\lim_{n \rightarrow \infty} \lambda_2 = 0$.

Summary A method has been presented which allows one to study rare events in random systems down to regions of *very low* probabilities. The basic idea is to represent the probability space by the phase space of a physical system held at temperature T . From the distribution of states, the original unbiased distribution can be obtained.

Here, the method is applied to the local sequence-alignment problem. A biologically relevant case is treated. The distribution of optimum alignment scores can be studied in regions where the probability is as small as 10^{-40} , and yet the deviations of the distribution from the theoretical prediction are visible. However, with

increasing sequence lengths, the distribution indeed approaches the Gumbel distribution. Hence, for very long sequences, when one is not interested in the deviation from the Gumbel form, the island method [18] is a more suitable tool to study sequence alignment. Please note that the island method is not a general purpose method like the technique presented here, but designed especially for local alignments.

Acknowledgements The author developed the idea for this method at the workshop “Statistical Physics of Biological Information” at the Institute for Theoretical Physics in Santa Barbara during discussions with P. Grassberger and E. Marinari. The author would like to thank A.P. Young and P. Grassberger for critically reading the manuscript and interesting discussions and A.P. Young also for various other support. The simulations were performed on a Beowulf Cluster at the Institut für Theoretische Physik of the Universität Magdeburg (Germany). Financial support was obtained from the DFG (Deutsche Forschungsgemeinschaft) under grant Ha 3169/1-1.

-
- [1] International Humane Genome Sequencing Consortium, *Nature* **409**, 860 (2001)
 - [2] S.M. Brown, *Bioinformatics*, (Eaton Publishing, Natick (MA) 2000)
 - [3] H.H. Rashidi and L.K. Buehler, *Bioinformatics Basics*, (CRC Press, Boca Raton (FL) 2000)
 - [4] S. Heinkoff and J.G. Heinkoff, *Proc. Natl. Acad. Sci. USA* **89**, 10915 (1992)
 - [5] O. Gotoh, *J. Mol. Biol.* **162**, 705 (1982)
 - [6] T.F. Smith, M.S. Waterman, and C. Burks, *Nucleic Acids Res.* **13**, 645 (1985)
 - [7] J.F. Collins, A.F.W. Coulson, and A. Lyall, *CABIOS* **4**, 67 (1988);
 - [8] R. Mott, *Bull. Math. Biol.* **54**, 59 (1992)
 - [9] M.S. Waterman and V. Vingron, *Proc. Natl. Acad. Sci. USA* **91**, 4625 (1994); *Stat. Sci.* **9**, 367 (1994)
 - [10] S.F. Altschul and W. Gish, *Methods in Enzymology* **266**, 460 (1996)
 - [11] E.J. Gumbel, *Statistics of Extremes*, (Columbia Univ. Press, New York 1958)
 - [12] B.D. Ripley, *Stochastic simulation*, (Wiley, New York 1987)
 - [13] B. Berg and T. Neuhaus, *Phys. Rev. Lett.* **68**, 9 (1992)
 - [14] P. Grassberger, *Phys. Rev. E* **56**, 3682 (1997)
 - [15] S. Kirkpatrick, C. D. Gelatt, and M. P. Vecchi, *Science* **220**, 671 (1983)
 - [16] S.B. Needleman and C.D. Wunsch, *J. Mol. Biol.* **48**, 443 (1970)
 - [17] T.F. Smith and M.S. Waterman, *J. Mol. Biol.* **147**, 195 (1981)
 - [18] S.F. Altschul, R. Bundschuh, R. Olsen, and T. Hwa, *Nucl. Acid Res.* **29**, 351 (2001)

- [19] S. Lipschutz and M. Lipson, *Probability*, (McGraw-Hill, New York 2000)
- [20] N. Metropolis, A. Rosenbluth, M. Rosenbluth, A. Teller, and E. Teller, J. Chem. Phys. **21**, 1087 (1953)
- [21] V. Kalashnikov, *Geometric Sums: Bounds for Rare Events with Applications*, (Kluwer Academic Publishers, Dordrecht 1997)
- [22] L.E. Reichl, *A Modern Course in Statistical Physics*, (Wiley, New York 1998)
- [23] E. Marinari and G. Parisi, Europhys. Lett. **19**, 451 (1992)
- [24] K. Hukushima and K. Nemoto, J. Phys. Soc. Jap. **65**, 1604 (1996)
- [25] Only when $\mathbf{x} = \mathbf{y}$ the highest possible score is achieved
- [26] A.B. Robinson and L.R. Robinson, Proc. Natl. Acad. Sci. USA, **88**, 8880 (1991)